CUMULATIVE AUTHOR INDEX
Annals of Entomology
Volume 94, 2001

A

Agrawal, A. A., 71
Akimoto, S.-I., 462
Albrecht, C., 702
Albrecht, C. P., 433
Albuquerque, G. S., 858
Allen, J. F., 654
Aluja, M., 268, 886
Alyokhin, A. V., 664
Andow, D. A., 563
Appel, A. G., 420
Armella, M. A., 16
Arroyave, J. A., 512
Asís, J. D., 809

B

Barrera, E., 174, 363
Belliure, B., 91
Bellotti, A., 512
Berkebile, D. R., 695
Birke, A., 268
Bishop, D. B., 33
Bosch, J., 244, 617
Bounfour, M., 400
Brailovsky, H., 174, 363
Bristow, C. M., 33
Brock, B. L., 732
Broers, D. A., 405
Broufas, G. D., 82
Brownie, C., 555
Buckner, J. S., 743, 761

C

Calkins, C. O., 405
Calvert, L. A., 512
Cameron, S. A., 851
Carey, J. R., 41
Cassis, G., 363
Chang, C. L., 433, 702
Chapco, W., 505
Charlton, R. E., 427
Chernoff, N., 555
Chu, C.-C., 743, 761
Clark, T. L., 877
Cohen, A. C., 146
Čokl, A., 471
Cole, B. J., 59
Cole, D., 851
Cônsoli, F. L., 97
Constantino, L. M., 185, 512
Cooley, J. R., 755
Costa, J. T., 480
Crist, T. O., 583
Cuervo, M., 512

D

de Brito, R. O. A., 333
Dejean, A., 449
De León, T., 197
Díaz-Fleischer, F., 268
Dobler, S., 1
dos Santos, P., 717
Douglas, L. J., 726

E

El-Shall, S. S. A., 702
Emoto, J., 467
Espinoza, A. M., 438
Etges, W. J., 16
Evans, E. W., 76, 605

F

Fadamiro, H. Y., 909
Feder, J. L., 902, 936
Filchak, K. E., 902
Foster, J. E., 877
Franz, N. M., 835
Fraser, L., 954
Freeman, T. P., 695, 743, 761
Frohlich, D., 512
Furniss, M. M., 21

G

Galante, E., 596
Gayubo, S. F., 809
Gilbert, L. E., 97
Göllner, U., 363
Gómez, J. M., 110
González-Megías, A., 110
Goyer, R. A., 948
Grant, J. F., 896
Greenberg, S. M., 566
Grimaldi, D., 10

H

Haight, K., 59
Hall, H. G., 670
Hall, J. P. W., 185, 631
Hamid, A. A., 545
Harvey, D. J., 631
Haymer, D. S., 726
Heed, W. B., 16
Heimpel, G. E., 909
Henneberry, T. J., 743, 761
Hernández-Ortiz, V., 268
Hirabayashi, K., 123
Holland, J., 1
Horton, D. R., 405
Hosoka·va, T., 750
Howard, R. W., 152, 707
Hunt, J. H., 555

I

Ibáñez-Bernal, S., 367
Ihringova, P., 179
Inoue, T., 545
Itioka, T., 545

J

Jacob, H. S., 605
Jameson, M. L., 866
Jones, R. W., 51

K

Kalaskar, A., 76
Kaliang, H., 545
Kambhampati, S., 732
Kato, M., 545
Katsoyannos, B. I., 41
Kemp, W. P., 244
Kidd, K. A., 219
Klompen, H., 10
Kopper, B. J., 427
Kouloussis, N. A., 41
Koveos, D. S., 82

Krecek, J., 530
Krombein, K. V., 809
Kruse, J. J., 166
Kurashima, R., 433, 702

L
Labandeira, C. C., 179
Lachaud, J.-P., 707
Leborgne, R., 490
Lee, C.-F., 347
Legaspi, B. C., Jr., 566
Leopold, R. A., 695
Leskey, T. C., 275
Lewis, C. N., 230
Lewis, M. L., 648
Lewis, R. R., 405
Litzenberger, G., 505
Liu, T.-X., 566
Locke, S., 1
Louque, R. W., 480
Lozada, N., 268
Lu, W., 64
Lundström, J. O., 576

M
Maeta, Y., 617
Malherbe, M. C., 449
Manfrin, M. H., 333
Marshall, D. C., 386
Matioli, S. R., 717
May-Itza, W. D. J., 540
McBrien, H. L., 471
McMichael, M. A., 670
McNeil, J. N., 226
Meinke, L. J., 877
Messing, R. H., 664
Messinger, S., 851
Metzler, E. H., 583
Michaud, J. P., 91
Miliczky, E. R., 405
Millar, J. G., 471
Miller, K. B., 520
Miyatake, T., 612
Mizota, K., 462
Mohamad, S. U., 545
Momose, K., 545
Moncayo, A. C., 503
Montgomery, M. E., 64
Mora, R., 438
Morn, M.-A., 771

N
Nagamitsu, T., 545
Naik, R. R., 686
Nakamoto, N., 123
Nakamuta, K., 917
Natwick, E. T., 743
Nelson, D. R., 743, 761
Newton, B. L., 116
Nice, C. C., 257
Nichols, B. J., 26
Nilsson, A. N., 769
Nishida, T., 550
Noda, A., 467
Noonan, G. R., 301

O
O'Brien, C. W., 835
O'Donnell, S., 555
O'Grady, P. M., 16
Oku, S., 550
Orivel, J., 449
Orr, D. B., 219

P
Papadopoulos, N. T., 41
Pasquet, A., 490
Pérez-Lachaud, G., 707
Peterson, M. A., 1
Phillips, P. K., 159
Piñero, J., 268
Platt, A. P., 654
Porter, S. D., 289
Powell, J. A., 677
Prinzing, A. J., 210
Prokopy, R. J., 275

Q
Quezada-Euan, J. J. G., 540

R
Ramaswamy, S. B., 427
Ramírez, S. R., 851
Rani, P. U., 917
Reid, W., 457
Retana, A., 438
Robacker, D. C., 954
Roehrdanz, R. L., 928
Roesch, K., 851
Roethele, J. B., 902, 936
Romero-Severson, J., 936
Rossi, A. M., 871
Ruberson, J. R., 116
Rust, R., 617

S
Sakai, S., 545
Sanborn, A. F., 159
Sappington, T. W., 415, 566
Satô, M., 347
Schaefer, C., 628
Schaefer, C. W., 298
Schäfer, M., 576
Scheffer, S. J., 648
Scheffrahn, R. H., 530
Schmitz, H., 686
Selivon, D., 592
Sene, F. M., 333
Sétamou, M., 566
Shapiro, A. M., 257
Shelly, T. E., 497
Shelton, T. G., 420
Sherriffs, M. F., 71
Shu, S., 427
Sites, R. W., 26
Sivinski, J., 886
Skoda, S. R., 877
Sousa, J. M., 681
Sowards, L. A., 686
Spence, J. R., 681
Sperling, F. A. H., 166
Steinmiller, B., 732
Stone, M. O., 686
Storozhenko, S. Y., 179
Strom, B. L., 948
Strong, D., 871
Summerville, K. S., 583
Suzuki, N., 750

T
Tanigoshi, L. K., 400
Tantalo, L., 1
Tatsuta, H., 462
Tauber, C. A., 197, 858
Tauber, M. J., 858
Taylor, O. M., 851
Thevenard, L., 490
Tisdale, R. A., 415
Tomlin, D. W., 686
Tormos, J., 809
Torres-Vila, L. M., 226

U
Uramoto, K., 717

V
Valdez, J. M., 251
Vander Meer, R. K., 289
Venner, S., 490
Verdú, J. R., 596
Villatoro, K., 866
Vinson, S. B., 97
Viviani, V. R., 129
Vrsansky, P., 179
Vulinec, K., 886

W
Wang, W. B., 695
Welbourn, W. C., 896
Whitfield, J. B., 230, 851
Wiernasz, D. C., 59
Wiggins, G. J., 896
Williams, B. L., 239
Willmott, K. R., 185
Wolf, K. W., 457
Wuellner, C. T., 97

Y
Yamada, S. M., 592
Yamane, S., 545
Yang, P., 664
Yang, P.-S., 347
Yeargan, K. V., 116
Yoshida, A., 467
Yumoto, T., 545

Z
Zeng, F., 146

CUMULATIVE KEYWORD INDEX
Annals of Entomology
Volume 94, 2001

A

abdominal marks,
Aporophyla nigra, 226
Hecatera bicolorata, 226
mating status, 226
Mythimna unipuncta, 226
Noctuid, 226
Acanthocephalini,
Bolivia, 174
Creidae, 174
Ichilocoris, 174
Peru, 174
acoustic behavior,
cicadas, 159
Diceroprocta delicata, 159
morphometrics, 159
systematics, 159
thermal adaptation, 159
acoustic response,
Chironomidae, 123
midge, 123
swarm, 123
wingbeat sound, 123
Acrosternum hilare,
mating behavior, 471
Nezara viridula, 471
pulse train, 471
stink bug, 471
substrate-borne vibrational communication, 471
adaptation,
digestive enzymes, 146
elastin, 146
Lygus hesperus, 146
plant bug, 146
Adelges tsugae,
development, 64
feeding, 64
hemlocks, 64
oviposition, 64
Scymnus sinuanodulus, 64
adhesive pad,
arboreal life, 449
arolium, 449
arolium gland, 449
Pachycondyla, 449
pretarsus, 449
adult feeding,
age-specific mortality, 563
European corn borer, 563
life expectancy, 563
sugar feeding, 563
Africanized honey bees,
allozyme, 540
Apis mellifera, 540
drone, 540
haplotype, 540
mating, 540
Agaporomorphus,
diving beetles, 520
Dytiscidae, 520
Neotropical, 520
phylogeny, 520
Agelaia panamaensis,
Agelaia yepocapa, 555
caste, 555
colony, 555
inquilines, 555
morphometrics, 555
Agelaia yepocapa,
Agelaia panamaensis, 555
caste, 555
colony, 555
inquilines, 555
morphometrics, 555
age-specific mortality,
adult feeding, 563
European corn borer, 563
life expectancy, 563
sugar feeding, 563
aggression,
alate, 289
fire ant, 289
nestmate recognition, 289
polygyny, 289
alate,
aggression, 289
fire ant, 289
nestmate recognition, 289
polygyny, 289

Aleyrodidae,
16S rDNA, 512
compound eye, 512
polymerase chain reaction, 512
random amplified polymorphic DNA, 512
whitefly, 512
alfalfa leafcutting bee,
development, 244
emergence, 244
Megachile rotundata, 244
mortality, 244
pollination, 244
alfalfa weevil,
biological control, 605
floral feeding, 605
host-habitat, 605
Hypera postica, 605
Medicago sativa, 605
allometry,
genitalia, 462
head, 462
lock-and-key hypothesis, 462
Lucanidae, 462
sexual selection hypothesis, 462
allozyme,
Africanized honey bees, 540
Apis mellifera, 540
drone, 540
haplotype, 540
mating, 540
allozymes, 1, 257
host race formation, 257
hybridization, 1
Lepidoptera, 257
mating, 1
mitochondrial DNA, 257
Mitoura, 257
morphology, 1
phylogeography, 257
Anadaptus,
Anisodactylus, 301
biogeography, 301
GIS (geographic information system), 301
Nearctic, 301
Neotropical, 301
Anastrepha,
fruit flies, 592
genetic marker, 592
ommatidia, 592
Anastrepha ludens,
Diptera, 251
histology, 251
morphology, 251
Tephritidae, 251
testis, 251
Angoumois grain moth,
chemical ecology, 152
cowpea weevil, 152
parasitoid, 152
Pteromalus cerealellae, 152
stored-product pest, 152
Anisodactylus,
Anadaptus, 301
biogeography, 301
GIS (geographic information system), 301
Nearctic, 301
Neotropical, 301
ants,
Cinara aphids, 33
Formica exsectoides, 33
homopteran community, 33
mutualism, 33
scale insects, 33
Aphis spiraecola,
Pseudodorus clavatus, 91
Syrphidae, 91
Toxoptera citricida, 91
Apis koschevnikovi,
Borneo, 545
general flowering, 545
migration, 545
numerical response, 545
Southeast Asian tropics, 545
Apis mellifera,
Africanized honey bees, 540
allozyme, 540
drone, 540
haplotype, 540
mating, 540
Aporophyla nigra,
abdominal marks, 226
Hecatera bicolorata, 226
mating status, 226
Mythimna unipuncta, 226
Noctuid, 226
Araneae,
orchards, 405
overwintering, 405
species diversity, 405

arboreal life,
adhesive pad, 449
arolium, 449
arolium gland, 449
Pachycondyla, 449
pretarsus, 449
Archipini,
Archips argyrospila host race, 166
COI, 166
pheromone race, 166
Tortricinae, 166
Archips argyrospila host race,
Archipini, 166
COI, 166
pheromone race, 166
Tortricinae, 166
Argasidae,
Carios, 10
fossil, 10
arolium,
adhesive pad, 449
arboreal life, 449
arolium gland, 449
Pachycondyla, 449
pretarsus, 449
arolium gland,
adhesive pad, 449
arboreal life, 449
arolium, 449
Pachycondyla, 449
pretarsus, 449
Ateuchini,
breeding patterns, 596
dung beetles, 596
larval description, 596
Scarabaeoidea, 596
taxonomy, 596
attractant,
Ceratitis capitata, 497
α-copaene, 497
mating behavior, 497
Australia,
Dicranocephalus, 363
Hemiptera, 363
Stenocephalidae, 363

B

bees,
behavioral characters, 617
molecular characters, 617
nesting phylogeny, 617
Osmia, 617
beet armyworm,
fecundity, 415
longevity, 415
oviposition, 415
Spodoptera exigua, 415
behavior, 210, 268
biophysical ecology, 210
canopy, 210
ecophysiology, 210
feeding, 268
habitat use, 210
mating system, 268
oviposition, 268
Rhagoletis turpiniae, 268
scale, 210
Tephritidae, 268
behavioral characters,
bees, 617
molecular characters, 617
nesting phylogeny, 617
Osmia, 617
Belize,
Bruchomyiinae, 367
new species, 367
Phlebotominae, 367
Psychodidae, 367
taxonomy, 367
biodiversity,
Braconidae, 230
grasslands, 230
parasitoids, 230
tallgrass prairies, 230
biogeography,
Anadaptus, 301
Anisodactylus, 301
GIS (geographic information system), 301
Nearctic, 301
Neotropical, 301
biological control, 76, 605
alfalfa weevil, 605
Coccinella septempunctata, 76
floral feeding, 605
foraging behavior, 76
Harmonia axyridis, 76
host-habitat, 605
Hypera postica, 76, 605
Medicago sativa, 605
predation, 76
bioluminescence,
fireflies, 129

habitats, 129
Lampyridae, 129
larvae, 129
biophysical ecology,
behavior, 210
canopy, 210
ecophysiology, 210
habitat use, 210
scale, 210
Bolivia,
Acanthocephalini, 174
Creidae, 174
Ichilocoris, 174
Peru, 174
boll weevil,
coevolution, 51
host plants, 51
phylogenetics, 51
Borneo,
Apis koschevnikovi, 545
general flowering, 545
migration, 545
numerical response, 545
Southeast Asian tropics, 545
Braconidae,
biodiversity, 230
grasslands, 230
parasitoids, 230
tallgrass prairies, 230
breeding patterns,
Ateuchini, 596
dung beetles, 596
larval description, 596
Scarabaeoidea, 596
taxonomy, 596
bristle distribution,
butterfly, 467
Pieris rapae, 467
wing margin, 467
brood extinction,
geographic distributions, 386
life cycles, 386
Magicicada, 386
periodical cicada, 386
phenotypic plasticity, 386
Bruchomyiinae,
Belize, 367
new species, 367
Phlebotominae, 367
Psychodidae, 367
taxonomy, 367
butterfly,
bristle distribution, 467
Pieris rapae, 467
wing margin, 467

C

cactus,
Drosophila, 16
Mexico, 16
systematics, 16
canopy,
behavior, 210
biophysical ecology, 210
ecophysiology, 210
habitat use, 210
scale, 210
capture thread length,
formulae, 490
orb webs, 490
spiders, 490
sticky silk, 490
web-building investment, 490
Carios,
Argasidae, 10
fossil, 10
caste,
Agelaia panamaensis, 555
Agelaia yepocapa, 555
colony, 555
inquilines, 555
morphometrics, 555
Ceratitis capitata, 41, 433, 497
attractant, 497
α-copaene, 497
host infestation, 41
mass rearing, 433
mating behavior, 497
meridic diet, 433
northern Greece, 41
population dynamics, 41
trapping, 41
chemical communication,
Group foraging, 480
insect sociality, 480
Sawfly, 480
Symphyta, 480
trail following, 480
chemical ecology,
Angoumois grain moth, 152
cowpea weevil, 152
parasitoid, 152
Pteromalus cerealellae, 152

stored-product pest, 152
Chironomidae,
acoustic response, 123
midge, 123
swarm, 123
wingbeat sound, 123
Chrysopidae,
larval morphology, 197
Mexico, 197
systematics, 197
trash-carriers, 197
Chysomelidae,
feeding preference, 110
host range, 110
specialization, 110
Timarcha, 110
cicadas,
acoustic behavior, 159
Diceroprocta delicata, 159
morphometrics, 159
systematics, 159
thermal adaptation, 159
Cinara aphids,
ants, 33
Formica exsectoides, 33
homopteran community, 33
mutualism, 33
scale insects, 33
clines,
conservation, 239
Speyeria, 239
taxonomy, 239
wing pattern, 239
CO_2 release patterns,
respiration, 420
subterranean termites, 420
water loss patterns, 420
Coccinella septempunctata,
biological control, 76
foraging behavior, 76
Harmonia axyridis, 76
Hypera postica, 76
predation, 76
coevolution,
boll weevil, 51
host plants, 51
phylogenetics, 51
COI,
Archipini, 166
Archips argyrospila host race, 166
pheromone race, 166
Tortricinae, 166
cold hardiness,
Euseius finlandicus, 82
Phytoseiidae, 82
supercooling, 82
Coleoptera,
Dendroctonus pseudotsugae barragani, 21
Mexico, 21
Scolytidae, 21
Colobura,
early stages, 185
hostplants, 185
Neotropical, 185
taxonomy, 185
colony,
Agelaia panamaensis, 555
Agelaia yepocapa, 555
caste, 555
inquilines, 555
morphometrics, 555
complementarity,
moth, 583
rarity, 583
scale dependence, 583
spatial extent, 583
species richness, 583
compound eye,
16S rDNA, 512
Aleyrodidae, 512
polymerase chain reaction, 512
random amplified polymorphic DNA, 512
whitefly, 512
Conotrachelus nenuphar,
host-plant volatiles, 275
pheromones, 275
conservation,
clines, 239
Speyeria, 239
taxonomy, 239
wing pattern, 239
conspecific suppression,
feeding behavior, 97
gas exchange, 97
larval morphology, 97
parasitoid–host interaction, 97
red imported fire ant, 97
consumption rates,
feeding growth indices, 566
host plants, 566

Spodoptera exigua, 566
α-copaene,
attractant, 497
Ceratitis capitata, 497
mating behavior, 497
cowpea weevil,
Angoumois grain moth, 152
chemical ecology, 152
parasitoid, 152
Pteromalus cerealellae, 152
stored-product pest, 152
Creidae,
Acanthocephalini, 174
Bolivia, 174
Ichilocoris, 174
Peru, 174
Cylas formicarius,
death mimicry, 612
sexual difference, 612
sweetpotato weevil, 612
thanatosis, 612

D

death mimicry,
Cylas formicarius, 612
sexual difference, 612
sweetpotato weevil, 612
thanatosis, 612
delphacid,
genitalia, 438
mouthparts, 438
sensilla, 438
Tagosodes orizicolus, 438
ultrastructure, 438
Dendroctonus pseudotsugae barragani,
Coleoptera, 21
Mexico, 21
Scolytidae, 21
development, 64, 244
Adelges tsugae, 64
alfalfa leafcutting bee, 244
emergence, 244
feeding, 64
hemlocks, 64
Megachile rotundata, 244
mortality, 244
oviposition, 64
pollination, 244
Scymnus sinuanodulus, 64
developmental rates,
Eotetranychus carpini borealis, 400
life history, 400
spider mites, 400
Tetranychidae, 400
Tetranychus urticae, 400
Diceroprocta delicata,
acoustic behavior, 159
cicadas, 159
morphometrics, 159
systematics, 159
thermal adaptation, 159
Dicranocephalus,
Australia, 363
Hemiptera, 363
Stenocephalidae, 363
digestive enzymes,
adaptation, 146
elastin, 146
Lygus hesperus, 146
plant bug, 146
Diptera,
Anastrepha ludens, 251
histology, 251
morphology, 251
Tephritidae, 251
testis, 251
distribution, 59, 530
Myrmecocystus, 59
Myrmecocystus mexicanus, 59
Neotropics, 530
new species, 530
Pogonomyrmex, 59
Pogonomyrmex occidentalis, 59
Procryptotermes, 530
spatial pattern, 59
West Indies, 530
Yucatan, 530
diversity,
functional groups, 576
mosquitoes, 576
wetlands, 576
diving beetles,
Agaporomorphus, 520
Dytiscidae, 520
Neotropical, 520
phylogeny, 520
drone,
Africanized honey bees, 540
allozyme, 540
Apis mellifera, 540
haplotype, 540
mating, 540

Drosophila,
cactus, 16
Mexico, 16
systematics, 16
Drosophila buzzatii cluster,
geographic variation., 333
mtDNA COI, 333
phylogeny, 333
dung beetles,
Ateuchini, 596
breeding patterns, 596
larval description, 596
Scarabaeoidea, 596
taxonomy, 596
Dytiscidae,
Agaporomorphus, 520
diving beetles, 520
Neotropical, 520
phylogeny, 520

E

early stages,
Colobura, 185
hostplants, 185
Neotropical, 185
taxonomy, 185
ecophysiology,
behavior, 210
biophysical ecology, 210
canopy, 210
habitat use, 210
scale, 210
egg structure,
life history, 26
Limnocoris lutzi, 26
lotic ecology, 26
Naucoridae, 26
taxonomy, 26
elastin,
adaptation, 146
digestive enzymes, 146
Lygus hesperus, 146
plant bug, 146
emergence,
alfalfa leafcutting bee, 244
development, 244
Megachile rotundata, 244
mortality, 244
pollination, 244
Eotetranychus carpini borealis,
developmental rates, 400
life history, 400
spider mites, 400
Tetranychidae, 400
Tetranychus urticae, 400
Eubrianacinae,
phylogenetics, 347
Psephenidae, 347
systematics, 347
Eurasia,
grasshoppers, 505
Melanoplinae, 505
mitochondrial DNA, 505
phylogeography, 505
Podismini, 505
European corn borer,
adult feeding, 563
age-specific mortality, 563
life expectancy, 563
sugar feeding, 563
Euseius finlandicus,
cold hardiness, 82
Phytoseiidae, 82
supercooling, 82

F

fecundity, 219, 415
beet armyworm, 415
host preference, 219
kudzu, 219
longevity, 415
oviposition, 415
preimaginal development, 219
Pseudoplusia includens, 219
soybean, 219
Spodoptera exigua, 415
feeding, 64, 268
Adelges tsugae, 64
behavior, 268
development, 64
hemlocks, 64
mating system, 268
oviposition, 64, 268
Rhagoletis turpiniae, 268
Scymnus sinuanodulus, 64
Tephritidae, 268
feeding behavior,
conspecific suppression, 97
gas exchange, 97
larval morphology, 97
parasitoid–host interaction, 97
red imported fire ant, 97
feeding growth indices,
consumption rates, 566

host plants, 566
Spodoptera exigua, 566
feeding preference,
Chysomelidae, 110
host range, 110
specialization, 110
Timarcha, 110
fire ant,
aggression, 289
alate, 289
nestmate recognition, 289
polygyny, 289
fireflies,
bioluminescence, 129
habitats, 129
Lampyridae, 129
larvae, 129
floral feeding,
alfalfa weevil, 605
biological control, 605
host-habitat, 605
Hypera postica, 605
Medicago sativa, 605
foraging behavior,
biological control, 76
Coccinella septempunctata, 76
Harmonia axyridis, 76
Hypera postica, 76
predation, 76
Formica exsectoides,
ants, 33
Cinara aphids, 33
homopteran community, 33
mutualism, 33
scale insects, 33
formulae,
capture thread length, 490
orb webs, 490
spiders, 490
sticky silk, 490
web-building investment, 490
fossil,
Argasidae, 10
Carios, 10
fruit flies,
Anastrepha, 592
genetic marker, 592
ommatidia, 592
functional groups,
diversity, 576
mosquitoes, 576
wetlands, 576

G

Galloisiana olgae,
Grylloblattodea, 179
new species, 179
paleobiology, 179
rock crawlers, 179
Ussuri River Basin, 179
gas exchange,
conspecific suppression, 97
feeding behavior, 97
larval morphology, 97
parasitoid–host interaction, 97
red imported fire ant, 97
general flowering,
Apis koschevnikovi, 545
Borneo, 545
migration, 545
numerical response, 545
Southeast Asian tropics, 545
generalist versus specialist,
induced defense, 71
plant-insect interactions, 71
Raphanus raphanistrum, 71
trade-offs, 71
genetic marker,
Anastrepha, 592
fruit flies, 592
ommatidia, 592
genitalia, 438, 462
allometry, 462
delphacid, 438
head, 462
lock-and-key hypothesis, 462
Lucanidae, 462
mouthparts, 438
sensilla, 438
sexual selection hypothesis, 462
Tagosodes orizicolus, 438
ultrastructure, 438
Geocoris punctipes,
life history, 116
photoperiod, 116
preoviposition period, 116
reproductive diapause, 116
geographic distributions,
brood extinction, 386
life cycles, 386
Magicicada, 386
periodical cicada, 386

phenotypic plasticity, 386
geographic variation.,
Drosophila buzzatii cluster, 333
mtDNA COI, 333
phylogeny, 333
GIS (geographic information system),
Anadaptus, 301
Anisodactylus, 301
biogeography, 301
Nearctic, 301
Neotropical, 301
grasshoppers,
Eurasia, 505
Melanoplinae, 505
mitochondrial DNA, 505
phylogeography, 505
Podismini, 505
grasslands,
biodiversity, 230
Braconidae, 230
parasitoids, 230
tallgrass prairies, 230
Group foraging,
chemical communication, 480
insect sociality, 480
Sawfly, 480
Symphyta, 480
trail following, 480
Grylloblattodea,
Galloisiana olgae, 179
new species, 179
paleobiology, 179
rock crawlers, 179
Ussuri River Basin, 179

H

habitat use,
behavior, 210
biophysical ecology, 210
canopy, 210
ecophysiology, 210
scale, 210
habitats,
bioluminescence, 129
fireflies, 129
Lampyridae, 129
larvae, 129
haplotype,
Africanized honey bees, 540
allozyme, 540
Apis mellifera, 540
drone, 540
mating, 540
Harmonia axyridis,
biological control, 76
Coccinella septempunctata, 76
foraging behavior, 76
Hypera postica, 76
predation, 76
head,
allometry, 462
genitalia, 462
lock-and-key hypothesis, 462
Lucanidae, 462
sexual selection hypothesis, 462
Hecatera bicolorata,
abdominal marks, 226
Aporophyla nigra, 226
mating status, 226
Mythimna unipuncta, 226
Noctuid, 226
Hemiptera,
Australia, 363
Dicranocephalus, 363
Stenocephalidae, 363
hemlocks,
Adelges tsugae, 64
development, 64
feeding, 64
oviposition, 64
Scymnus sinuanodulus, 64
histology,
Anastrepha ludens, 251
Diptera, 251
morphology, 251
Tephritidae, 251
testis, 251
homopteran community,
ants, 33
Cinara aphids, 33
Formica exsectoides, 33
mutualism, 33
scale insects, 33
host infestation,
Ceratitis capitata, 41
northern Greece, 41
population dynamics, 41
trapping, 41
host plants, 51, 566
boll weevil, 51
coevolution, 51
consumption rates, 566
feeding growth indices, 566

phylogenetics, 51
Spodoptera exigua, 566
host preference,
fecundity, 219
kudzu, 219
preimaginal development, 219
Pseudoplusia includens, 219
soybean, 219
host race formation,
allozymes, 257
Lepidoptera, 257
mitochondrial DNA, 257
Mitoura, 257
phylogeography, 257
host range,
Chysomelidae, 110
feeding preference, 110
specialization, 110
Timarcha, 110
host-habitat,
alfalfa weevil, 605
biological control, 605
floral feeding, 605
Hypera postica, 605
Medicago sativa, 605
host-plant volatiles,
Conotrachelus nenuphar, 275
pheromones, 275
hostplants,
Colobura, 185
early stages, 185
Neotropical, 185
taxonomy, 185
hybridization,
allozymes, 1
mating, 1
morphology, 1
Hypera postica, 76, 605
alfalfa weevil, 605
biological control, 76, 605
Coccinella septempunctata, 76
floral feeding, 605
foraging behavior, 76
Harmonia axyridis, 76
host-habitat, 605
Medicago sativa, 605
predation, 76

I

Ichilocoris,
Acanthocephalini, 174
Bolivia, 174
Creidae, 174
Peru, 174
induced defense,
generalist versus specialist, 71
plant-insect interactions, 71
Raphanus raphanistrum, 71
trade-offs, 71
inquilines,
Agelaia panamaensis, 555
Agelaia yepocapa, 555
caste, 555
colony, 555
morphometrics, 555
insect sociality,
chemical communication, 480
Group foraging, 480
Sawfly, 480
Symphyta, 480
trail following, 480
integumental glands,
preying behavior, 457
Reduviidae, 457
setae, 457
sundew hairs, 457
Zelus longipes, 457

J

juvenile hormone,
monandry, 427
Nymphalidae, 427
oocyte maturation, 427
reproductive strategy, 427
Speyeria idalia, 427

K

kudzu,
fecundity, 219
host preference, 219
preimaginal development, 219
Pseudoplusia includens, 219
soybean, 219

L

Lampyridae,
bioluminescence, 129
fireflies, 129
habitats, 129
larvae, 129
larvae,
bioluminescence, 129
fireflies, 129

habitats, 129
Lampyridae, 129
larval description,
Ateuchini, 596
breeding patterns, 596
dung beetles, 596
Scarabaeoidea, 596
taxonomy, 596
larval morphology, 97, 197
Chrysopidae, 197
conspecific suppression, 97
feeding behavior, 97
gas exchange, 97
Mexico, 197
parasitoid–host interaction, 97
red imported fire ant, 97
systematics, 197
trash-carriers, 197
Lepidoptera,
allozymes, 257
host race formation, 257
mitochondrial DNA, 257
Mitoura, 257
phylogeography, 257
life cycles,
brood extinction, 386
geographic distributions, 386
Magicicada, 386
periodical cicada, 386
phenotypic plasticity, 386
life expectancy,
adult feeding, 563
age-specific mortality, 563
European corn borer, 563
sugar feeding, 563
life history, 26, 116, 400
developmental rates, 400
egg structure, 26
Eotetranychus carpini borealis, 400
Geocoris punctipes, 116
Limnocoris lutzi, 26
lotic ecology, 26
Naucoridae, 26
photoperiod, 116
preoviposition period, 116
reproductive diapause, 116
spider mites, 400
taxonomy, 26
Tetranychidae, 400
Tetranychus urticae, 400
Limnocoris lutzi,
egg structure, 26
life history, 26
lotic ecology, 26
Naucoridae, 26
taxonomy, 26
local mate competition,
sex ratio, 550
single-sex brood, 550
Trypoxylon malaisei, 550
lock-and-key hypothesis,
allometry, 462
genitalia, 462
head, 462
Lucanidae, 462
sexual selection hypothesis, 462
longevity,
beet armyworm, 415
fecundity, 415
oviposition, 415
Spodoptera exigua, 415
lotic ecology,
egg structure, 26
life history, 26
Limnocoris lutzi, 26
Naucoridae, 26
taxonomy, 26
Lucanidae,
allometry, 462
genitalia, 462
head, 462
lock-and-key hypothesis, 462
sexual selection hypothesis, 462
Lygus hesperus,
adaptation, 146
digestive enzymes, 146
elastin, 146
plant bug, 146

M

Magicicada,
brood extinction, 386
geographic distributions, 386
life cycles, 386
periodical cicada, 386
phenotypic plasticity, 386
mass rearing,
Ceratitis capitata, 433
meridic diet, 433
mating, 1, 540
Africanized honey bees, 540
allozyme, 540

allozymes, 1
Apis mellifera, 540
drone, 540
haplotype, 540
hybridization, 1
morphology, 1
mating behavior, 471, 497
Acrosternum hilare, 471
attractant, 497
Ceratitis capitata, 497
α-copaene, 497
Nezara viridula, 471
pulse train, 471
stink bug, 471
substrate-borne vibrational communication, 471
mating status,
abdominal marks, 226
Aporophyla nigra, 226
Hecatera bicolorata, 226
Mythimna unipuncta, 226
Noctuid, 226
mating system,
behavior, 268
feeding, 268
oviposition, 268
Rhagoletis turpiniae, 268
Tephritidae, 268
Medicago sativa,
alfalfa weevil, 605
biological control, 605
floral feeding, 605
host-habitat, 605
Hypera postica, 605
Megachile rotundata,
alfalfa leafcutting bee, 244
development, 244
emergence, 244
mortality, 244
pollination, 244
Melanoplinae,
Eurasia, 505
grasshoppers, 505
mitochondrial DNA, 505
phylogeography, 505
Podismini, 505
meridic diet,
Ceratitis capitata, 433
mass rearing, 433
Mexico, 16, 21, 197
cactus, 16
Chrysopidae, 197
Coleoptera, 21
Dendroctonus pseudotsugae barragani, 21
Drosophila, 16
larval morphology, 197
Scolytidae, 21
systematics, 16, 197
trash-carriers, 197
midge,
acoustic response, 123
Chironomidae, 123
swarm, 123
wingbeat sound, 123
migration,
Apis koschevnikovi, 545
Borneo, 545
general flowering, 545
numerical response, 545
Southeast Asian tropics, 545
mitochondrial DNA, 257, 505
allozymes, 257
Eurasia, 505
grasshoppers, 505
host race formation, 257
Lepidoptera, 257
Melanoplinae, 505
Mitoura, 257
phylogeography, 257, 505
Podismini, 505
Mitoura,
allozymes, 257
host race formation, 257
Lepidoptera, 257
mitochondrial DNA, 257
phylogeography, 257
molecular characters,
bees, 617
behavioral characters, 617
nesting phylogeny, 617
Osmia, 617
monandry,
juvenile hormone, 427
Nymphalidae, 427
oocyte maturation, 427
reproductive strategy, 427
Speyeria idalia, 427
morphology, 1, 251
allozymes, 1
Anastrepha ludens, 251
Diptera, 251

histology, 251
hybridization, 1
mating, 1
Tephritidae, 251
testis, 251
morphometrics, 159, 555
acoustic behavior, 159
Agelaia panamaensis, 555
Agelaia yepocapa, 555
caste, 555
cicadas, 159
colony, 555
Diceroprocta delicata, 159
inquilines, 555
systematics, 159
thermal adaptation, 159
mortality,
alfalfa leafcutting bee, 244
development, 244
emergence, 244
Megachile rotundata, 244
pollination, 244
mosquitoes,
diversity, 576
functional groups, 576
wetlands, 576
moth,
complementarity, 583
rarity, 583
scale dependence, 583
spatial extent, 583
species richness, 583
mouthparts,
delphacid, 438
genitalia, 438
sensilla, 438
Tagosodes orizicolus, 438
ultrastructure, 438
mtDNA COI,
Drosophila buzzatii cluster, 333
geographic variation., 333
phylogeny, 333
mutualism,
ants, 33
Cinara aphids, 33
Formica exsectoides, 33
homopteran community, 33
scale insects, 33
Myrmecocystus,
distribution, 59
Myrmecocystus mexicanus, 59
Pogonomyrmex, 59
Pogonomyrmex occidentalis, 59
spatial pattern, 59
Myrmecocystus mexicanus,
distribution, 59
Myrmecocystus, 59
Pogonomyrmex, 59
Pogonomyrmex occidentalis, 59
spatial pattern, 59
Mythimna unipuncta,
abdominal marks, 226
Aporophyla nigra, 226
Hecatera bicolorata, 226
mating status, 226
Noctuid, 226

N

Naucoridae,
egg structure, 26
life history, 26
Limnocoris lutzi, 26
lotic ecology, 26
taxonomy, 26
Nearctic,
Anadaptus, 301
Anisodactylus, 301
biogeography, 301
GIS (geographic information system), 301
Neotropical, 301
Neotropical, 185, 301, 520
Agaporomorphus, 520
Anadaptus, 301
Anisodactylus, 301
biogeography, 301
Colobura, 185
diving beetles, 520
Dytiscidae, 520
early stages, 185
GIS (geographic information system), 301
hostplants, 185
Nearctic, 301
phylogeny, 520
taxonomy, 185
Neotropics,
distribution, 530
new species, 530
Procryptotermes, 530
West Indies, 530
Yucatan, 530
nesting phylogeny,

bees, 617
behavioral characters, 617
molecular characters, 617
Osmia, 617
nestmate recognition,
aggression, 289
alate, 289
fire ant, 289
polygyny, 289
new species, 179, 367, 530
Belize, 367
Bruchomyiinae, 367
distribution, 530
Galloisiana olgae, 179
Grylloblattodea, 179
Neotropics, 530
paleobiology, 179
Phlebotominae, 367
Procryptotermes, 530
Psychodidae, 367
rock crawlers, 179
taxonomy, 367
Ussuri River Basin, 179
West Indies, 530
Yucatan, 530
Nezara viridula,
Acrosternum hilare, 471
mating behavior, 471
pulse train, 471
stink bug, 471
substrate-borne vibrational communication, 471
Noctuid,
abdominal marks, 226
Aporophyla nigra, 226
Hecatera bicolorata, 226
mating status, 226
Mythimna unipuncta, 226
northern Greece,
Ceratitis capitata, 41
host infestation, 41
population dynamics, 41
trapping, 41
numerical response,
Apis koschevnikovi, 545
Borneo, 545
general flowering, 545
migration, 545
Southeast Asian tropics, 545
Nymphalidae,
juvenile hormone, 427
monandry, 427
oocyte maturation, 427
reproductive strategy, 427
Speyeria idalia, 427

O

ommatidia,
Anastrepha, 592
fruit flies, 592
genetic marker, 592
oocyte maturation,
juvenile hormone, 427
monandry, 427
Nymphalidae, 427
reproductive strategy, 427
Speyeria idalia, 427
orb webs,
capture thread length, 490
formulae, 490
spiders, 490
sticky silk, 490
web-building investment, 490
orchards,
Araneae, 405
overwintering, 405
species diversity, 405
Osmia,
bees, 617
behavioral characters, 617
molecular characters, 617
nesting phylogeny, 617
overwintering,
Araneae, 405
orchards, 405
species diversity, 405
oviposition, 64, 268, 415
Adelges tsugae, 64
beet armyworm, 415
behavior, 268
development, 64
fecundity, 415
feeding, 64, 268
hemlocks, 64
longevity, 415
mating system, 268
Rhagoletis turpiniae, 268
Scymnus sinuanodulus, 64
Spodoptera exigua, 415
Tephritidae, 268

P

Pachycondyla,
adhesive pad, 449
arboreal life, 449
arolium, 449
arolium gland, 449
pretarsus, 449
paleobiology,
Galloisiana olgae, 179
Grylloblattodea, 179
new species, 179
rock crawlers, 179
Ussuri River Basin, 179
parasitoid,
Angoumois grain moth, 152
chemical ecology, 152
cowpea weevil, 152
Pteromalus cerealellae, 152
stored-product pest, 152
parasitoid–host interaction,
conspecific suppression, 97
feeding behavior, 97
gas exchange, 97
larval morphology, 97
red imported fire ant, 97
parasitoids,
biodiversity, 230
Braconidae, 230
grasslands, 230
tallgrass prairies, 230
periodical cicada,
brood extinction, 386
geographic distributions, 386
life cycles, 386
Magicicada, 386
phenotypic plasticity, 386
Peru,
Acanthocephalini, 174
Bolivia, 174
Creidae, 174
Ichilocoris, 174
phenotypic plasticity,
brood extinction, 386
geographic distributions, 386
life cycles, 386
Magicicada, 386
periodical cicada, 386
pheromone race,
Archipini, 166
Archips argyrospila host race, 166
COI, 166
Tortricinae, 166
pheromones,
Conotrachelus nenuphar, 275
host-plant volatiles, 275
Phlebotominae,
Belize, 367
Bruchomyiinae, 367
new species, 367
Psychodidae, 367
taxonomy, 367
photoperiod,
Geocoris punctipes, 116
life history, 116
preoviposition period, 116
reproductive diapause, 116
phylogenetics, 51, 347
boll weevil, 51
coevolution, 51
Eubrianacinae, 347
host plants, 51
Psephenidae, 347
systematics, 347
phylogeny, 333, 520
Agaporomorphus, 520
diving beetles, 520
Drosophila buzzatii cluster, 333
Dytiscidae, 520
geographic variation., 333
mtDNA COI, 333
Neotropical, 520
phylogeography, 257, 505
allozymes, 257
Eurasia, 505
grasshoppers, 505
host race formation, 257
Lepidoptera, 257
Melanoplinae, 505
mitochondrial DNA, 257, 505
Mitoura, 257
Podismini, 505
Phytoseiidae,
cold hardiness, 82
Euseius finlandicus, 82
supercooling, 82
Pieris rapae,
bristle distribution, 467
butterfly, 467
wing margin, 467
plant bug,
adaptation, 146
digestive enzymes, 146

elastin, 146
Lygus hesperus, 146
plant-insect interactions,
generalist versus specialist, 71
induced defense, 71
Raphanus raphanistrum, 71
trade-offs, 71
Podismini,
Eurasia, 505
grasshoppers, 505
Melanoplinae, 505
mitochondrial DNA, 505
phylogeography, 505
Pogonomyrmex,
distribution, 59
Myrmecocystus, 59
Myrmecocystus mexicanus, 59
Pogonomyrmex occidentalis, 59
spatial pattern, 59
Pogonomyrmex occidentalis,
distribution, 59
Myrmecocystus, 59
Myrmecocystus mexicanus, 59
Pogonomyrmex, 59
spatial pattern, 59
pollination,
alfalfa leafcutting bee, 244
development, 244
emergence, 244
Megachile rotundata, 244
mortality, 244
polygyny,
aggression, 289
alate, 289
fire ant, 289
nestmate recognition, 289
polymerase chain reaction,
16S rDNA, 512
Aleyrodidae, 512
compound eye, 512
random amplified polymorphic DNA, 512
whitefly, 512
population dynamics,
Ceratitis capitata, 41
host infestation, 41
northern Greece, 41
trapping, 41
predation,
biological control, 76
Coccinella septempunctata, 76
foraging behavior, 76
Harmonia axyridis, 76
Hypera postica, 76
preimaginal development,
fecundity, 219
host preference, 219
kudzu, 219
Pseudoplusia includens, 219
soybean, 219
preoviposition period,
Geocoris punctipes, 116
life history, 116
photoperiod, 116
reproductive diapause, 116
pretarsus,
adhesive pad, 449
arboreal life, 449
arolium, 449
arolium gland, 449
Pachycondyla, 449
preying behavior,
integumental glands, 457
Reduviidae, 457
setae, 457
sundew hairs, 457
Zelus longipes, 457
Procryptotermes,
distribution, 530
Neotropics, 530
new species, 530
West Indies, 530
Yucatan, 530
Psephenidae,
Eubrianacinae, 347
phylogenetics, 347
systematics, 347
Pseudodorus clavatus,
Aphis spiraecola, 91
Syrphidae, 91
Toxoptera citricida, 91
Pseudoplusia includens,
fecundity, 219
host preference, 219
kudzu, 219
preimaginal development, 219
soybean, 219
Psychodidae,
Belize, 367
Bruchomyiinae, 367
new species, 367

Phlebotominae, 367
taxonomy, 367
Pteromalus cerealellae,
Angoumois grain moth, 152
chemical ecology, 152
cowpea weevil, 152
parasitoid, 152
stored-product pest, 152
pulse train,
Acrosternum hilare, 471
mating behavior, 471
Nezara viridula, 471
stink bug, 471
substrate-borne vibrational communication, 471

R

random amplified polymorphic DNA,
16S rDNA, 512
Aleyrodidae, 512
compound eye, 512
polymerase chain reaction, 512
whitefly, 512
Raphanus raphanistrum,
generalist versus specialist, 71
induced defense, 71
plant-insect interactions, 71
trade-offs, 71
rarity,
complementarity, 583
moth, 583
scale dependence, 583
spatial extent, 583
species richness, 583
red imported fire ant,
conspecific suppression, 97
feeding behavior, 97
gas exchange, 97
larval morphology, 97
parasitoid–host interaction, 97
Reduviidae,
integumental glands, 457
preying behavior, 457
setae, 457
sundew hairs, 457
Zelus longipes, 457
reproductive diapause,
Geocoris punctipes, 116
life history, 116
photoperiod, 116
preoviposition period, 116
reproductive strategy,
juvenile hormone, 427
monandry, 427
Nymphalidae, 427
oocyte maturation, 427
Speyeria idalia, 427
respiration,
CO_2 release patterns, 420
subterranean termites, 420
water loss patterns, 420
Rhagoletis turpiniae,
behavior, 268
feeding, 268
mating system, 268
oviposition, 268
Tephritidae, 268
rock crawlers,
Galloisiana olgae, 179
Grylloblattodea, 179
new species, 179
paleobiology, 179
Ussuri River Basin, 179

S

Sawfly,
chemical communication, 480
Group foraging, 480
insect sociality, 480
Symphyta, 480
trail following, 480
scale,
behavior, 210
biophysical ecology, 210
canopy, 210
ecophysiology, 210
habitat use, 210
scale dependence,
complementarity, 583
moth, 583
rarity, 583
spatial extent, 583
species richness, 583
scale insects,
ants, 33
Cinara aphids, 33
Formica exsectoides, 33
homopteran community, 33
mutualism, 33
Scarabaeoidea,
Ateuchini, 596
breeding patterns, 596
dung beetles, 596

larval description, 596
taxonomy, 596
Scolytidae,
Coleoptera, 21
Dendroctonus pseudotsugae barragani, 21
Mexico, 21
Scymnus sinuanodulus,
Adelges tsugae, 64
development, 64
feeding, 64
hemlocks, 64
oviposition, 64
sensilla,
delphacid, 438
genitalia, 438
mouthparts, 438
Tagosodes orizicolus, 438
ultrastructure, 438
setae,
integumental glands, 457
preying behavior, 457
Reduviidae, 457
sundew hairs, 457
Zelus longipes, 457
sex ratio,
local mate competition, 550
single-sex brood, 550
Trypoxylon malaisei, 550
sexual difference,
Cylas formicarius, 612
death mimicry, 612
sweetpotato weevil, 612
thanatosis, 612
sexual selection hypothesis,
allometry, 462
genitalia, 462
head, 462
lock-and-key hypothesis, 462
Lucanidae, 462
single-sex brood,
local mate competition, 550
sex ratio, 550
Trypoxylon malaisei, 550
Southeast Asian tropics,
Apis koschevnikovi, 545
Borneo, 545
general flowering, 545
migration, 545
numerical response, 545
soybean,
fecundity, 219
host preference, 219
kudzu, 219
preimaginal development, 219
Pseudoplusia includens, 219
spatial extent,
complementarity, 583
moth, 583
rarity, 583
scale dependence, 583
species richness, 583
spatial pattern,
distribution, 59
Myrmecocystus, 59
Myrmecocystus mexicanus, 59
Pogonomyrmex, 59
Pogonomyrmex occidentalis, 59
specialization,
Chysomelidae, 110
feeding preference, 110
host range, 110
Timarcha, 110
species diversity,
Araneae, 405
orchards, 405
overwintering, 405
species richness,
complementarity, 583
moth, 583
rarity, 583
scale dependence, 583
spatial extent, 583
Speyeria,
clines, 239
conservation, 239
taxonomy, 239
wing pattern, 239
Speyeria idalia,
juvenile hormone, 427
monandry, 427
Nymphalidae, 427
oocyte maturation, 427
reproductive strategy, 427
spider mites,
developmental rates, 400
Eotetranychus carpini borealis, 400
life history, 400
Tetranychidae, 400
Tetranychus urticae, 400

spiders,
capture thread length, 490
formulae, 490
orb webs, 490
sticky silk, 490
web-building investment, 490
Spodoptera exigua, 415, 566
beet armyworm, 415
consumption rates, 566
fecundity, 415
feeding growth indices, 566
host plants, 566
longevity, 415
oviposition, 415
Stenocephalidae,
Australia, 363
Dicranocephalus, 363
Hemiptera, 363
sticky silk,
capture thread length, 490
formulae, 490
orb webs, 490
spiders, 490
web-building investment, 490
stink bug,
Acrosternum hilare, 471
mating behavior, 471
Nezara viridula, 471
pulse train, 471
substrate-borne vibrational communication, 471
stored-product pest,
Angoumois grain moth, 152
chemical ecology, 152
cowpea weevil, 152
parasitoid, 152
Pteromalus cerealellae, 152
substrate-borne vibrational communication,
Acrosternum hilare, 471
mating behavior, 471
Nezara viridula, 471
pulse train, 471
stink bug, 471
subterranean termites,
CO_2 release patterns, 420
respiration, 420
water loss patterns, 420
sugar feeding,
adult feeding, 563
age-specific mortality, 563
European corn borer, 563
life expectancy, 563
sundew hairs,
integumental glands, 457
preying behavior, 457
Reduviidae, 457
setae, 457
Zelus longipes, 457
supercooling,
cold hardiness, 82
Euseius finlandicus, 82
Phytoseiidae, 82
swarm,
acoustic response, 123
Chironomidae, 123
midge, 123
wingbeat sound, 123
sweetpotato weevil,
Cylas formicarius, 612
death mimicry, 612
sexual difference, 612
thanatosis, 612
Symphyta,
chemical communication, 480
Group foraging, 480
insect sociality, 480
Sawfly, 480
trail following, 480
Syrphidae,
Aphis spiraecola, 91
Pseudodorus clavatus, 91
Toxoptera citricida, 91
systematics, 16, 159, 197, 347
acoustic behavior, 159
cactus, 16
Chrysopidae, 197
cicadas, 159
Diceroprocta delicata, 159
Drosophila, 16
Eubrianacinae, 347
larval morphology, 197
Mexico, 16, 197
morphometrics, 159
phylogenetics, 347
Psephenidae, 347
thermal adaptation, 159
trash-carriers, 197

T

Tagosodes orizicolus,
delphacid, 438
genitalia, 438

mouthparts, 438
sensilla, 438
ultrastructure, 438
tallgrass prairies,
biodiversity, 230
Braconidae, 230
grasslands, 230
parasitoids, 230
taxonomy, 26, 185, 239, 367, 596
Ateuchini, 596
Belize, 367
breeding patterns, 596
Bruchomyiinae, 367
clines, 239
Colobura, 185
conservation, 239
dung beetles, 596
early stages, 185
egg structure, 26
hostplants, 185
larval description, 596
life history, 26
Limnocoris lutzi, 26
lotic ecology, 26
Naucoridae, 26
Neotropical, 185
new species, 367
Phlebotominae, 367
Psychodidae, 367
Scarabaeoidea, 596
Speyeria, 239
wing pattern, 239
Tephritidae, 251, 268
Anastrepha ludens, 251
behavior, 268
Diptera, 251
feeding, 268
histology, 251
mating system, 268
morphology, 251
oviposition, 268
Rhagoletis turpiniae, 268
testis, 251
testis,
Anastrepha ludens, 251
Diptera, 251
histology, 251
morphology, 251
Tephritidae, 251
Tetranychidae,
developmental rates, 400
Eotetranychus carpini borealis, 400
life history, 400
spider mites, 400
Tetranychus urticae, 400
Tetranychus urticae,
developmental rates, 400
Eotetranychus carpini borealis, 400
life history, 400
spider mites, 400
Tetranychidae, 400
thanatosis,
Cylas formicarius, 612
death mimicry, 612
sexual difference, 612
sweetpotato weevil, 612
thermal adaptation,
acoustic behavior, 159
cicadas, 159
Diceroprocta delicata, 159
morphometrics, 159
systematics, 159
Timarcha,
Chysomelidae, 110
feeding preference, 110
host range, 110
specialization, 110
Tortricinae,
Archipini, 166
Archips argyrospila host race, 166
COI, 166
pheromone race, 166
Toxoptera citricida,
Aphis spiraecola, 91
Pseudodorus clavatus, 91
Syrphidae, 91
trade-offs,
generalist versus specialist, 71
induced defense, 71
plant-insect interactions, 71
Raphanus raphanistrum, 71
trail following,
chemical communication, 480
Group foraging, 480
insect sociality, 480
Sawfly, 480
Symphyta, 480
trapping,
Ceratitis capitata, 41
host infestation, 41
northern Greece, 41
population dynamics, 41

trash-carriers,
Chrysopidae, 197
larval morphology, 197
Mexico, 197
systematics, 197
Trypoxylon malaisei,
local mate competition, 550
sex ratio, 550
single-sex brood, 550

U

ultrastructure,
delphacid, 438
genitalia, 438
mouthparts, 438
sensilla, 438
Tagosodes orizicolus, 438
Ussuri River Basin,
Galloisiana olgae, 179
Grylloblattodea, 179
new species, 179
paleobiology, 179
rock crawlers, 179

W

water loss patterns,
CO_2 release patterns, 420
respiration, 420
subterranean termites, 420
web-building investment,
capture thread length, 490
formulae, 490
orb webs, 490
spiders, 490
sticky silk, 490
West Indies,
distribution, 530
Neotropics, 530
new species, 530
Procryptotermes, 530
Yucatan, 530
wetlands,
diversity, 576
functional groups, 576
mosquitoes, 576
whitefly,
16S rDNA, 512
Aleyrodidae, 512
compound eye, 512
polymerase chain reaction, 512
random amplified polymorphic DNA, 512
wing margin,
bristle distribution, 467
butterfly, 467
Pieris rapae, 467
wing pattern,
clines, 239
conservation, 239
Speyeria, 239
taxonomy, 239
wingbeat sound,
acoustic response, 123
Chironomidae, 123
midge, 123
swarm, 123

Y

Yucatan,
distribution, 530
Neotropics, 530
new species, 530
Procryptotermes, 530
West Indies, 530

Z

Zelus longipes,
integumental glands, 457
preying behavior, 457
Reduviidae, 457
setae, 457
sundew hairs, 457